



Pyola[®] Insect Spray

**Fast acting—
kills on contact**

ACTIVE INGREDIENTS:	By Wt.
Pyrethrins	0.5%
Canola Oil	89.5%
OTHER INGREDIENTS	10.0%
TOTAL	100.0%

This product contains 0.859 lbs. of canola oil and 0.005 lbs. of pyrethrins per pint.

**KEEP OUT OF REACH
OF CHILDREN**

CAUTION

See back panels for first aid
and directions for use.

NET CONTENTS: 16 Fl. Oz.

64415

EPA Reg. No. 67702-5-56872, EPA Est. No. 50591-GA-01

DORMANT USE

Kills egg stages of pests present in dormant season and scale insects, mites, and mealybugs on dormant shrubs (e.g., roses), evergreens, fruit trees and shade trees. Apply a 2% solution. Thoroughly wet the dormant plants or trees. Do not apply more than 4 fl.oz. of concentrate (400 fl.oz. of 1% solution or 200 fl.oz. of 2% solution) per 1000 square feet.

Precautionary Statements

Hazards to Humans and Domestic Animals:

CAUTION: Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

User Safety Recommendations

- Users should wash hands with plenty of soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards: This product is toxic to aquatic organisms including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Notice To Buyer To the extent consistent with applicable law, Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. To the extent consistent with applicable law, this warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller. To the extent consistent with applicable law, the Seller shall not be liable for any such use. To the extent consistent with applicable law, the Seller makes no other warranties, either expressed or implied.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep this product in its tightly closed original container when not in use. Store in a cool, dry, well-ventilated (preferably locked) area that is inaccessible to children and animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call the poison control center at 1-800-222-1222.



Made with Pyrol™ Insecticide a trademark of W. Neudorff GmbH KG

Distributed by Gardens Alive!, Inc.
706 W. Cumberland St. Lewisburg, OH 45338
Phone: 513-354-1483 www.GardensAlive.com



Pyrola[®] Insect Spray

- Easy to use.
- Fast acting-kills on contact
- Contains pyrethrins and canola oil botanical insecticides
- Dormant and growing season insect spray to kill adelgids, aphids, armored scale, beetle larvae, canker worms, caterpillars, fungus gnats, lace bugs, leafhoppers, leafminers, leafrollers, mealybugs, mites, plant bugs, psyllids, scale insects, spider mites, thrips, webworms and whiteflies.
- May be used on edible fruits and vegetables up to and including day of harvest.

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency information concerning this product, call the poison control center at 1-800-222-1222.

KEEP OUT OF REACH OF CHILDREN CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals:

CAUTION: Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

User Safety Recommendations

- Users should wash hands with plenty of soap and water before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards: This product is toxic to aquatic organisms including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or

surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. **Only for use in home gardens and home greenhouses.**

USE RESTRICTIONS

- Do not wet plants to point of runoff or drip.
- Do not allow adults, children, or pets to enter the treated area until sprays have dried.
- Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift.
- Remove pets, birds, and cover fish aquariums before spraying.
- Do not apply within 21 days of a sulfur application.
- Do not use Pyrola Insect Spray when temperatures are above 90° F.
- Do not apply directly to or near water, storm drains, or drainage ditches. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) or apply when heavy rain is expected. Rinse applicator over lawn or garden area only.
- Do not apply more than 10 times per season.
- Do not reapply within 3 days except under extreme pest pressure.
- In case of extreme pest pressure do not re-apply within 24 hours.
- Not for use in outdoor residential misting systems.
- Do not make applications during rain.
- Do not water the treated area to the point of run-off.

Peel back tab for Storage and Disposal Instructions.



Made with Pyrol[™] Insecticide a trademark of W. Neudorff GmbH KG

BATCH CODE:

Distributed by Gardens Alive[®], Inc.
706 W. Cumberland St. Lewisburg, OH 45338
Phone: 513-354-1483 www.GardensAlive.com



1 92665 04954 5

Phytotoxicity Note:

Oil removes the glaucous (blue) bloom from such evergreens as Colorado blue spruce and Koster spruce. Use a 1% solution on oil sensitive plants such as: Chamaecyparis, Cryptomeria, Japanese Holly, Red Cedar, and Smoke Tree. Use care on blooms and buds. Test spray a portion of the plant and wait several days to determine if burn occurs before spraying larger areas. Some plants are sensitive to this product. If in doubt, test-spray a small portion of the plant and wait several days to determine if burn occurs before spraying larger areas.

Read entire label before use. Use only as directed. Shake well before each use. Thoroughly wet all parts of the plant. Repeat treatment in 1-2 weeks if necessary. May be used on edible fruits and vegetables up to and including day of harvest.

APPLICATION DIRECTIONS**INSTRUCTIONS FOR MIXING**

For a 1% solution, mix 2 teaspoons of concentrate with 1 quart (32 fl.oz.) water, or 1 fl.oz. concentrate with 3 quarts (96 fl.oz.) water. For a 2% solution, mix 4 teaspoons of concentrate with 1 quart (32 fl.oz.) water, or 2 fl.oz. concentrate with 3 quarts (96 fl.oz.) water. Shake well before each use and stir or shake diluted spray frequently during use.

PESTS/SITES AND APPLICATION**INSTRUCTIONS**

Pyola Insect Spray will kill: Adelgids, Ants (including Foraging Ants, Sweet-Feeding Ants, Pavement Ants, Moisture Ants, Little Black Ants, Argentine Ants, Crazy Ants, Acrobat Ants, Big-headed Ants, Cornfield Ants, Ghost Ants, Odorous House Ants, Thief Ants, Texas Leafcutting Ants, Field Ants, and other Common Ants) (except Fire Ants, Harvester Ants, Carpenter Ants, and Pharaoh Ants), Aphids (including Green Peach Aphids), Armyworms (including Beet Army Worms, Western Yellow Striped Armyworms), Bagworms, Beetles (Colorado potato beetle, flea beetle, Japanese beetle, asparagus beetle, Blister Beetles, 12-spotted Cucumber Beetles, Cucumber Beetle, Darkling Beetles (lesser mealworm), Saw-toothed Grain Beetles, Elm Leaf Beetles, Bean Beetles, Mexican Bean Beetles), Blow Flies, Borers (including European Corn Borer, Branch and Twig Borers, Shot hole borers and Pacific flathead borers), Box Elder Bugs, Cabbageworms (including Cross-striped Cabbageworm and Imported Cabbageworm), Cankerworms, Caterpillars (Gypsy Moth Caterpillars, Tent Caterpillar, Eastern Tent Caterpillar, Forest Tent Caterpillar and Tomato Fruitworm Caterpillar), Corn Earworms, Crane Flies, Crickets, Diamondback Moth Larvae, Earwigs, Fireworms, Fungus Gnats, Fruit Flies (including Cherry Fruit Fly), Glassy Winged Sharpshooter,

Grape Leaf Skeletonizer, Grasshoppers, Green Fruit Worm, Harlequin Bugs, Hornworms (including Tomato Hornworm), Katydid, Lace Bugs, Leafhoppers (including Grape Leafhopper, and Potato Leafhoppers), Leafminers, Leafrollers (including Fruittree Leafrollers), Leaf-tiers, Loopers (including Cabbage Looper), Lygus, Maggots (including Apple Maggot, Cabbage Maggots, and Onion Maggot), Mealybugs (including Citrus Mealybugs, Grape Mealybugs and Vine Mealybugs), Non-biting Midges, Mite-pedes, Mites (including Clover Mites, European Red Mite and Spider Mites), Moths (including Artichoke Plume Moth, Codling Moth, Diamondback Moth, European Pine Tip Moth, Gypsy Moths, Indian Meal Moth, Mediterranean Flour Moth), Mushroom Flies, Navel Orangeworm, Plant Bugs (including Leaf footed Plant Bug, Tarnished Plant Bug), Psyllids (including Pear Psyllids), Sawfly Larvae (pear and rose slugs), Scale insects (including Soft Scale and Armored Scale), Silverfish, Skippers, Sowbugs, Spittlebugs, Squash Bugs, Stink Bugs, Thrips (including Greenhouse Thrips), Vinegar Flies, Webworms, Weevils (including Boll Weevil, Carrot Weevil, Clover Weevil, Rice Weevil, Pepper Weevil), Whitefly and Phylloxera.

Apply at a rate of 1% or 2% solution. For soft-bodied pests such as aphids and mites, a 1% solution is sufficient. Using hand-held trigger sprayers, hand sprayers, or backpack sprayers, thoroughly spray insect pests with diluted product until plants are wet. Do not wet plants to point of runoff or drip. Repeat treatment in 1-2 weeks, if needed.

HOME GARDEN USE

For use outdoors on fruit and nut trees (apples, cherries, peaches, pears, nuts); flowering foliage plants; corn; melons; tomatoes; vegetables (beans, cabbage, cucurbits, peas, potatoes); figs; small fruits (grapes, strawberry, raspberry); citrus; ornamental and shade trees (birch, evergreens, holly, oak). Do not apply more than 4 fl.oz. of concentrate (400 fl.oz. of 1% solution or 200 fl.oz. of 2% solution) per 1000 square feet.

DIRECTIONS FOR GREENHOUSE USE

For use in home greenhouses on flowering, foliage plants; melons; tomatoes; vegetables (beans, cabbage, cucurbits, peas, potatoes); small fruits (grapes, strawberry, raspberry). Do not apply more than 4 fl.oz. of concentrate (400 fl.oz. of 1% solution or 200 fl.oz. of 2% solution) per 1000 square feet.